

§ 524.1044f

signs. If conditions persists or increases, discontinue use and consult veterinarian.

[48 FR 41157, Sept. 14, 1983, as amended 52 FR 7833, Mar. 13, 1987]

§ 524.1044f Gentamicin sulfate, betamethasone valerate topical spray.

(a) *Specifications.* Each milliliter of spray contains gentamicin sulfate equivalent to 0.57 milligram of gentamicin base and betamethasone valerate equivalent to 0.284 milligram of betamethasone.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs in the treatment of infected superficial lesions caused by bacteria sensitive to gentamicin.

(2) For the treatment of infected superficial lesions, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. Hold bottle upright 3 to 6 inches from the lesion and depress the sprayer head twice. One actuation of the sprayer delivers 0.7 milliliter of the spray. The drug should be administered with two spray actuations 2 to 4 times daily for 7 days.

(3) If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. The antibiotic susceptibility of the pathogenic organism should be determined prior to use of this preparation. Administration of recommended doses beyond 7 days may result in delayed wound healing. Animals treated longer than 7 days should be monitored closely.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 740, Jan. 7, 1985, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 10220, Mar. 6, 1997]

§ 524.1044g Gentamicin sulfate, betamethasone valerate, clotrimazole ointment.

(a) *Specifications.* Each gram (g) of ointment contains gentamicin sulfate equivalent to 3 milligrams (mg) gentamicin base, betamethasone valerate equivalent to 1 mg betamethasone, and 10 mg clotrimazole.

21 CFR Ch. I (4–1–05 Edition)

(b) *Sponsors.* See sponsors in § 510.600(c) of this chapter for uses as in paragraph (c) of this section.

(1) No. 000061 for use of 7.5- or 15-gram (g) tubes, 12.5-, 30-, or 215-g bottles.

(2) No. 051259 for use of 7.5- or 15-g tubes, 10-, or 25-g bottles.

(3) No. 059130 for use of 10-, 20-, or 215-g bottles.

(c) *Conditions of use in dogs—*(1) *Amount.* Instill ointment twice daily into the ear canal. Therapy should continue for 7 consecutive days.

(i) From 7.5- or 15-g tubes, 10-, 12.5-, 25-, or 30-g bottles: 4 drops for dogs weighing less than 30 pounds (lb) or 8 drops for dogs weighing 30 lb or more.

(ii) From 20- or 215-g bottles: 2 drops for dogs weighing less than 30 lb or 4 drops for dogs weighing 30 lb or more.

(2) *Indications for use.* For the treatment of acute and chronic canine otitis externa associated with yeast (*Malassezia pachydermatis*, formerly *Pityrosporum canis*) and/or bacteria susceptible to gentamicin.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 38973, July 21, 1993, as amended at 63 FR 31932, June 11, 1998; 68 FR 42970, July 21, 2003; 70 FR 8291, Feb. 18, 2005]

§ 524.1044h Gentamicin sulfate, mometasone furoate, clotrimazole otic suspension.

(a) *Specifications.* Each gram contains gentamicin sulfate, United States Pharmacopeia (USP) equivalent to 3 milligram (mg) gentamicin base, mometasone furoate monohydrate equivalent to 1 mg mometasone, and 10 mg clotrimazole, USP.

(b) *Sponsor.* See No. 000061 in § 510.6(c) of this chapter.

(c) *Conditions of use—Dogs—*(1) *Amount.* For dogs weighing less than 30 pounds (lb), instill 4 drops from the 5- and 30-gram (g) bottle into the ear canal (2 drops from the 215-g bottle) or, for dogs weighing 30 lb or more, instill 8 drops from the 5- and 30-g bottle into the ear canal (4 drops from the 215-g bottle), once or twice daily for 7 days.

(2) *Indications for use.* For the treatment of otitis externa caused by susceptible strains of yeast (*Malassezia pachydermatis*) and bacteria (*Pseudomonas* spp. [including *P.*